



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Patterns of care in recurrent glioblastoma in Switzerland: a multicentre national approach based on diagnostic nodes

Hundsberger, T ; Hottinger, A F ; Roelcke, U ; Roth, P ; Migliorini, D ; Dietrich, P Y ; Conen, K ; Pesce, G ; Hermann, E ; Pica, A ; Gross, M W ; Brügge, D ; Plasswilm, L ; Weller, M ; Putora, P M

Abstract: Despite moderate improvements in outcome of glioblastoma after first-line treatment with chemoradiation recent clinical trials failed to improve the prognosis of recurrent glioblastoma. In the absence of a standard of care we aimed to investigate institutional treatment strategies to identify similarities and differences in the pattern of care for recurrent glioblastoma. We investigated re-treatment criteria and therapeutic pathways for recurrent glioblastoma of eight neuro-oncology centres in Switzerland having an established multidisciplinary tumour-board conference. Decision algorithms, differences and consensus were analysed using the objective consensus methodology. A total of 16 different treatment recommendations were identified based on combinations of eight different decision criteria. The set of criteria implemented as well as the set of treatments offered was different in each centre. For specific situations, up to 6 different treatment recommendations were provided by the eight centres. The only wide-range consensus identified was to offer best supportive care to unfit patients. A majority recommendation was identified for non-operable large early recurrence with unmethylated MGMT promoter status in the fit patients: here bevacizumab was offered. In fit patients with late recurrent non-operable MGMT promoter methylated glioblastoma temozolomide was recommended by most. No other majority recommendations were present. In the absence of strong evidence we identified few consensus recommendations in the treatment of recurrent glioblastoma. This contrasts the limited availability of single drugs and treatment modalities. Clinical situations of greatest heterogeneity may be suitable to be addressed in clinical trials and second opinion referrals are likely to yield diverging recommendations.

DOI: <https://doi.org/10.1007/s11060-015-1957-0>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-115210>

Journal Article

Accepted Version

Originally published at:

Hundsberger, T; Hottinger, A F; Roelcke, U; Roth, P; Migliorini, D; Dietrich, P Y; Conen, K; Pesce, G; Hermann, E; Pica, A; Gross, M W; Brügge, D; Plasswilm, L; Weller, M; Putora, P M (2016). Patterns of care in recurrent glioblastoma in Switzerland: a multicentre national approach based on diagnostic nodes. *Journal of Neuro-Oncology*, 126(1):175-183.

DOI: <https://doi.org/10.1007/s11060-015-1957-0>

Patterns of Care in Recurrent Glioblastoma in Switzerland - a multicentre national approach based on diagnostic nodes

Hundsberger T¹, Hottinger AF², Roelcke U³, Roth P⁴, Migliorini D⁵, Dietrich PY⁵, Conen K⁶, Pesce G⁷, Hermann E⁸, Pica A⁸, Gross MW⁹, Brügge D¹⁰, Plasswilm L¹⁰, Weller M⁴, Putora PM¹⁰

1 Dept. of Neurology and Dept. of Haematology/Oncology, Cantonal Hospital St. Gallen, CH

2 Dept. of Clinical Neurosciences & Oncology, University Hospital, Lausanne, CH

3 Brain Tumour Centre, Cantonal Hospital Aarau, CH

4 Dept. of Neurology, and Brain Tumor Centre, University Hospital, Zurich, CH

5 Dept. of Oncology, University Hospital Geneva, CH

6 Dept. of Oncology, University Hospital Basel, CH

7 Dept. of Radiation Oncology, Oncology Institute of Southern Switzerland, Bellinzona-Lugano, CH

8 Dept. of Radiation Oncology, University Hospital Bern, CH

9 Dept. of Radiation Oncology, University Hospital Basel, CH

10 Dept. of Radiation Oncology, Cantonal Hospital St. Gallen, CH

On behalf of the Swiss Group for Clinical Cancer Research (SAKK) Central Nervous System working group.

Correspondence:

Thomas Hundsberger, MD

Department of Neurology and

Department of Haematology/Oncology

Cantonal Hospital

Rorschacherstr. 95

9007 St. Gallen

Switzerland

Phone: 0041 71 494 3095

Fax: 0041 71 494 2895

Email: thomas.hundsberger@kssg.ch

Abstract

Background: Despite moderate improvements in outcome of glioblastoma after first-line treatment with chemoradiation recent clinical trials failed to improve the prognosis of recurrent glioblastoma. In the absence of a standard of care we aimed to investigate institutional treatment strategies to identify similarities and differences in the pattern of care for recurrent glioblastoma.

Methods: We investigated re-treatment criteria and therapeutic pathways for recurrent glioblastoma of 8 neuro-oncology centres in Switzerland having an established multidisciplinary tumour-board conference. Decision algorithms, differences and consensus were analysed using the objective consensus methodology.

Results: A total of 16 different treatment recommendations were identified based on combinations of 8 different decision criteria. The set of criteria implemented as well as the set of treatments offered was different in each centre. For specific situations, up to 6 different treatment recommendations were provided by the 8 centres. The only wide-range consensus identified was to offer best supportive care to unfit patients. A majority recommendation was identified for non-operable large early recurrence with unmethylated MGMT promoter status in the fit patients: here bevacizumab was offered. In fit patients with late recurrent non-operable MGMT promoter methylated glioblastoma temozolomide was recommended by most. No other majority recommendations were present.

Conclusion: In the absence of strong evidence we identified few consensus recommendations in the treatment of recurrent glioblastoma. This contrasts the limited availability of single drugs and treatment modalities. Clinical situations of greatest heterogeneity may be suitable to be addressed in clinical trials and second opinion referrals are likely to yield diverging recommendations.

Keywords: recurrent glioblastoma, diagnostic nodes, objective consensus, decision making, re-treatment criteria

Introduction

Glioblastoma is the most common and devastating primary brain tumour with an annual incidence of about three per 100000 [1] persons. It is associated with great morbidity and early mortality. Even after multimodal treatment with temozolomide chemoradiation (RT/TMZ→TMZ) progression-free survival (PFS) and overall survival (OS) remain poor with 6.9 and 14.2 months, respectively [2]. Recently, four large phase III trials failed to improve outcome after first-line therapy in terms of OS [3-6]. Hence, effective salvage treatment of recurrent glioblastoma remains a demanding problem in neuro-oncology.

The oral alkylating agent TMZ is widely used for the treatment of recurrent glioblastoma (TMZ re-challenge) [7-11]. However, the optimal dosing schedule and the minimal time interval from first line TMZ are unknown. This topic was investigated in a single prospective phase II trial using a metronomic TMZ schedule [9]. Patients with a tumour recurrence at least three months after RT/TMZ or two months after completion of RT/TMZ and six maintenance cycles of TMZ reached modest PFS rates after six months (PFS-6) of 27.3% and 35.7%, respectively. Recently, two dose-intensified TMZ schedules have been prospectively investigated at first relapse after at least two cycles of maintenance TMZ after RT/TMZ to avoid interference with pseudoprogression [11]. None of the regimes were superior with a median time to treatment failure of 1.8 months [Arm A: one week on (120 mg/m² per day)/one week off] and 2.0 months [Arm B: three weeks on (80 mg/m² per day)/ one week off], respectively. However, methylation of the MGMT promoter was a strong positive prognostic factor of PFS-6 (methylated 39.7% versus unmethylated 6.9%) [11]. Small targeted drugs like enzastaurin or cediranib failed to show benefit in recent phase III trials when compared to lomustine in recurrent glioblastoma [12, 13]. Therefore, lomustine has advanced to a widely used alkylating drug either alone or in combination whenever TMZ is not considered to be appropriate for the treatment of recurrent glioblastoma [14, 15]. In contrast to the European Union, bevacizumab, an anti-angiogenic agent targeting VEGF-A, has been approved in selected countries (i.e. North America, Switzerland) for the treatment of recurrent glioblastoma based on uncontrolled phase II studies [16, 17].

Due to the limited efficacy of second-line systemic treatment, focal strategies with re-operation and re-irradiation have also been introduced as salvage treatments. However, little evidence for these therapies is available as most studies investigated only small or retrospective patient cohorts [18-25]. Furthermore, availability of stereotactic radiotherapy or radiosurgery and financial resources of health care systems account for significant differences in use between countries and even national regions.

As effective treatment options are scarce and controlled trials are rare for recurrent glioblastoma, therapeutic decisions are mostly based on low level evidence [26]. Beyond evidence, experience (eminence-based medicine) may provide additional guidance. Possibly, further information can be extracted from the community, for example as patterns of care studies [27]. Patterns of care studies often rely on specific scenarios which are presented to participants, these are then analysed; the limitation being that the answers are restricted to only these specific scenarios. When the collected information is appropriately formatted [28], e.g. decision trees are collected, patterns of practice can

be analysed for multiple combinations of parameters [29]. In the setting of low evidence for salvage treatment we aimed to collect treatment algorithms from Swiss neuro-oncological centres which derive their treatment recommendations from multidisciplinary tumour boards. The collected information was used to perform a comprehensive patterns care and patterns of algorithms analysis for the treatment of recurrent glioblastoma.

Materials and Methods

Based on specific parameters (characteristics of the patient and the disease) recommendations can be defined. The concept of diagnostic nodes (Dodes) has been developed [28, 29] to allow for a cross-comparison of recommendations in decision-tree format and has been applied in a clinical setting [30, 31]. Dodes are organized into decision trees using pre-defined categories. Standardized nomenclature of parameters and recommendations is a prerequisite for automated comparisons and an unbiased evaluation. Based on Dodes and the objective consensus methodology [29], consensus and heterogeneity were analysed (Fig. 1).

Hospitals with an interdisciplinary neuro-oncology unit including at least full-time service of neurosurgery, radiotherapy and medical neuro-oncology with a dedicated neuro-oncology tumour board being represented in the Swiss Group for clinical cancer research (SAKK) were asked to participate in the study. Treatment recommendations for recurrent glioblastoma were collected without specifications in formatting. Free-text, Microsoft PowerPoint slides and hand-drawn diagrams were converted without changing content into treatment algorithms and after several bilateral exchanges defined as final by 1st of January 2015. These were again discussed among the participating centres. Two centres took patient preference explicitly into account; these were related to the decision for active treatment versus best supportive care (BSC). Despite these criteria being important for daily practice, they were eliminated for the analysis as they do not rely on a tumour board decision and of course should apply to every clinical decision. Similar criteria were fused into new comprehensive categories to simplify their usage and enable cross-comparability (i.e. the criteria fitness based upon performance status, co-morbidities and age). The resulting data were presented to the participants and minor corrections were applied. The interpretation by all participants resulted in the final discussion.

Results

In total, 8 Swiss centres (Aarau, Basel, Bellinzona, Bern, Geneva, Lausanne, St.Gallen, Zurich) participated and provided written or schematic information of their interdisciplinary local treatment strategy for recurrent glioblastoma. The original treatment algorithms included a total of 23 re-treatment criteria and a list of 16 treatment options (Table 1, Fig. 2).

Processing and simplification

As the considerable variability of the re-treatment criteria lead to an exponential rise in possible combinations, we simplified and integrated similar criteria with minor differences (Table 1). For example, the various thresholds of Karnofsky performance scale (KPS), age, Eastern cooperative oncology group performance status (ECOG) and the presence of co-morbidities were summarized as “fit” or “unfit”. Additionally, tumours were summarized as “resectable” or “non-resectable” according to their occurrence in eloquent areas and non-diffuse/localized distribution. Need for steroid use and symptomatic recurrent glioblastoma were merged to only one category. After simplification the trees were analysed with the so defined 8 essential criteria. All centres used resectability, fitness and time of recurrence as re-treatment criteria. Five centres (A, C-E, H) used the MGMT promoter status. However, we gathered no information which method and threshold for MGMT promoter methylation status was used in clinical practise. Four centres used tumour size (B, C, G, H) and unifocality (C, E, F, H). Only three and two centres, respectively, took symptoms (F, G, H) and complete resection of enhancing tumour (CRET) (A, B) into account.

Re-treatment criteria

All centres dichotomised their decision to actively treat recurrent glioblastoma with anti-tumour strategies by using the performance status (PS). Generally, “unfit” patients with a low PS were referred to BSC without offering active oncological treatment. The KPS and the ECOG were both reported. However, the lowest threshold for active anti-tumour re-treatment varied considerably (KPS from 50 to 90; ECOG 0-1) (Table 1). The same was true for age (between 50 and 75 years). However, age was only reported as a re-treatment criterion in 2 out of 8 centres.

Re-treatment options

Five treatments were offered against recurrent glioblastoma in at least 4 centres (Fig. 2). These were the combination of re-operation with either TMZ or bevacizumab, monotherapy with either TMZ or bevacizumab and BSC. Interestingly, the time required to consider TMZ re-challenge ranged from 2 to 6 months. Four centres additionally relied on the MGMT promoter methylation status to indicate an alkylator-based systemic treatment (TMZ or lomustine). Various re-treatment modalities and schemes were reported among the centres (Fig. 2). These included re-operation, re-irradiation, chemotherapy or immunotherapy, either as a monotherapy or as a multimodal treatment. For example, therapy varied by the combination of anti-neoplastic agents (bevacizumab, lomustine, temozolomide, bevacizumab/lomustine given in standard dosing), the schedule of re-irradiation (40 Gy in 1.66 fractions, 42 Gy in 2.66 fractions, 35 Gy in 3.5 fractions, 20 Gy in 2.0 fractions), re-irradiation with and without TMZ or bevacizumab and the method of re-irradiation (stereotactic radiotherapy (SRT), 3D-conformal or intensity modulated radiotherapy). Furthermore, criteria like the localisation and distribution of the tumour (local/distant recurrence; single/multiple/diffuse recurrence; operable/not-operable) were also taken into account (Table 1).

Consensus treatment strategies

Fig. 3 displays a comprehensive and condensed view of the re-treatment recommendations for recurrent glioblastoma where a majority recommendation was present (5 out of 8 centres; 63%). Seven out of 8 re-treatment criteria are implemented (time of recurrence, operability, size of the tumour, symptoms, fitness, MGMT promoter methylation and unifocality). The CRET criterion, which was only used in 2 centres (A, B), was not relevant to reach a majority treatment recommendation. Strong consensus was generally achieved in unfit patients. In all but one scenario 6 out of 8 centres recommended BSC instead of an anti-tumour treatment (fit=no). In contrast, in fit patients majority consensus was rare (fit=yes). Five out of 8 centres would treat fit patients with bevacizumab when a large non-operable early recurrent glioblastoma with an unmethylated MGMT promoter is present (* in Fig. 3). Additionally, TMZ would be the preferred recommendation (# in Fig. 3) in fit patients with a late recurrent, non-unifocal, non-operable and large glioblastoma with a methylated MGMT promoter with (5 out of 8 centres, 63%) or without clinical symptoms (6 out of 8 centres, 75%).

Discussion

In the absence of a standard of care for recurrent glioblastoma the aim of the present study was to investigate applied treatment strategies in 8 neuro-oncology centres in Switzerland. As we anticipated great heterogeneity we further aimed to identify clinical criteria which lead to individualized treatment decisions and to check if these criteria overlap between different centres. Our expectations in this regard were met as we identified various treatment modalities and multiple schedules as well as a plethora of various criteria for the treatment of recurrent glioblastoma. Based on these results, it is very unlikely that patients with recurrent glioblastoma obtain the same treatment recommendation

twice if they seek a second opinion in Switzerland. On the one hand this is due to individualized treatments for a given patient and takes patient- and centre-specific factors into account (i.e. availability of certain treatment options). On the other hand different recommendation can cause a decision dilemma for the patient.

Original parameters could be processed and reformatted with the help of a standardized procedure (Dodes). This method enables the identification of the most common treatment recommendations for any specific parameter combination [29]. The process was feasible and produced the most commonly agreed upon treatment recommendations derived from heterogeneous recommendations. The data presented here might therefore serve as one basis to develop a nationwide treatment guideline. Moreover, areas of controversy or low overlap can be used to address clinically important questions for future clinical trials.

All centres used the PS to decide about the indication of a tumour specific therapy. PS was measured with the KPS and ECOG which are well known instruments of daily oncological practice [14]. Additionally, age was a relevant criterion to guide treatment in some centres. PS and age are established prognostic factors for first-line as well as for second-line treatments [14, 32-36]. Recent studies set a threshold of KPS 60% (corresponding to ECOG 2) as a prerequisite for tumour specific treatment interventions [4, 5, 37, 38] as inferior PS is associated with increased side effects from any intervention and lowers the chance of clinical benefit [14, 36, 39]. The greatest consensus in our analysis was identified in unfit patients corresponding to a low PS. These patients would be mostly referred to BSC as the strategy of choice independent of operability, MGMT promoter methylation or time of recurrence, among other criteria (Fig. 3). Accordingly, this approach represents a robust consensus recommendation. This result is not trivial, as three out of eight centres did not actively claimed BSC as a primary re-treatment modality even after several feed-back rounds (Fig. 2). However, we are confident that a low PS would lead to the implementation of BSC without additional tumour specific treatment in daily practise in all centres. This issue might reflect a limitation of the study that not all applied treatment strategies are specifically enumerated by the participants.

Despite the negative prognostic impact of mental and cognitive decline [39] none of the centres reported mental status or a formal cognitive testing (i.e. MiniMental Status Exam) as an aid to decide about treatment for recurrent glioblastoma. It is tempting to speculate that the physicians felt they integrate all dimensions of neurological function in clinical decision making and do not need these scales for whatever reason (i.e. time consuming procedure). Co-morbidities and their formal investigation were also rarely taken into account (i.e. Charlson co-morbidity score) and none of the centres reported usage of a structured geriatric assessment [40-43]. Hence, the issue of fitness for re-treatment in recurrent glioblastoma mostly relies on a subjective perspective of the physician.

Various strategies of re-operation, re-irradiation, systemic therapy and multimodal treatments were reported [14]. Major level of consensus was reached for TMZ re-challenge in lesions with a methylated MGMT promoter even in the absence of a controlled trial at the time of investigation. Only recently, the DIRECTOR and the BELOB trial demonstrated the prognostic value of the MGMT-

promoter methylation for PFS-6 and PFS-9, respectively, with significantly better outcome in recurrent glioblastoma with a methylated MGMT promoter [11, 15]. However, the optimal time span between first-line and second-line TMZ at recurrence still remains elusive.

The anti-angiogenic agent bevacizumab is approved for the treatment of recurrent glioblastoma in selected countries including Switzerland. Somewhat surprisingly, only in fit, non-operable patients with a large lesion and an unmethylated MGMT promoter a majority of centres (63%) would offer bevacizumab. No consensus was obvious in all other occasions, despite approval and its steroid-sparing properties. The reason for this low overlap might be related to the absence of a controlled clinical trial for the use of bevacizumab in recurrent glioblastoma or the recently reported failure in the first-line treatment setting [4, 38]. Moreover, conflicting results about a negative impact on neuro-cognition may also hamper widespread use of this compound [44, 45].

Re-irradiation alone or in combination with systemic treatment did not reach any consensus. In addition to the paucity of controlled clinical trial this might reflect heterogeneity of radiation dose and technique. Despite the low efficacy of re-irradiation as a monotherapy observed in a controlled phase II trial with PFS-6 rates of 3.8% [24] a second radiation either alone or in combination with systemic therapy still remained a salvage treatment in five out of eight centres. However, this applies only in a selected patient population with good prognostic criteria. Of note, the same is true for re-operation which is mostly recommended in combination with other treatments in 7 out of 8 centres. Additionally, a controlled clinical trial demonstrating efficacy of this approach is also lacking. However, positive prognostic criteria (KPS > 80, tumour size \leq 50 ml, non-eloquent region) for the estimation of prolonged postoperative survival after re-operation have been reported [46]. In this regard it has to be mentioned that local treatment strategies are driven by the experience of the local physicians, the availability to diverse radiation techniques and the attitude of neuro-surgeons to offer glioblastoma resection at recurrence.

Conclusion

The neuro-oncologist may be equally soothed or unsettled by these results. For certain clinical scenarios, 8 centres provided 6 different treatment recommendations. This sheds an interesting light on the value and risk of obtaining a second opinion in this context. Despite experience and access to published literature, the interpretation and clinical implementation of this was very different among Swiss neuro-oncology centres.

The application of a decision tree analysis (objective consensus) was able to identify decision criteria relevant in clinical practice across centres. Future trials and guidelines should take these criteria into account to ease the transition of trial results and recommendations into clinical neuro-oncology.

Disclosure

1 TH, LP received honaria for advisory board lecture and travel grants from Roche and MSD. PR
2 received honoraria for advisory board activities or lectures from MSD, Roche, Novartis and Molecular
3 Partners. MW has received research grants from Acceleron, Actelion, Alpinia Institute, Bayer, Isarna,
4 MSD, Merck & Co, Novocure, PIQUR and Roche and honoraria for lectures or advisory board
5 participation or consulting from Celldex, Immunocellular, Isarna, Magforce, MSD, Merck & Co,
6 Northwest Biotherapeutics, Novocure, Pfizer, Roche and Teva. AFH, UR, DM, PYD, KC, EH, AP,
7 MWG, DB disclosed no conflict of interest.
8
9

10 11 12 *Ethical standards*

13 No specific patient data was used in this project.
14
15
16

17 18 *Funding*

19 The Department of Radiation Oncology (PMP) at the Cantonal hospital St. Gallen received a grant
20 from Roche, Switzerland to perform this analysis.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1

Overview of original re-treatment criteria, assessment procedures and the derived standardized criteria used for comparison.

Fig. 1 Analysis by the objective consensus method

Example of an early recurrent, small and resectable glioblastoma harbouring a methylated MGMT (mMGMT) promoter. For fit patients heterogeneous treatment options were recommended among the centres (A-H), whereas in unfit patients 6 out of 8 centres (A, B, E, G, H, F) suggested BSC and only two centres (C, D) recommended alternative treatments. Bev, bevacizumab; Lom, lomustine; OP, operation; TMZ, temozolomide; BSC, best supportive care; SRS, stereotactic radiosurgery; &, combined treatment; +, sequential treatment.

Fig. 2

Implemented treatment strategies among the centres A-H (cross, reported; white, not reported); OP, operation; SRS, stereotactic radiotherapy; Bev, bevacizumab; TMZ, temozolomide; RT, radiotherapy; LOM, lomustine; BSC, best supportive care.

Fig. 3

The most common recommendations tree reveals areas of consensus (at least 5 out of 8 centres, >63%) for patients who are unfit (A-F). The most common recommendation is best supportive care (BSC). Only on 3 clinical scenarios either bevacizumab (BEV, *) or temozolomide (TMZ, #) (B,E,F) is recommended by the majority of centres depending on the presence of symptoms. Only 7 out of 8 re-treatment criteria (all but the CRET criterion, see Fig. 3) were relevant for achieving a majority recommendation.

References

1. Ostrom QT, Bauchet L, Davis FG, Deltour I, Fisher JL, Langer CE, Pekmezci M, Schwartzbaum JA, Turner MC, Walsh KM (2014) The epidemiology of glioma in adults: a "state of the science" review. *Neuro-oncology* 16: 896-913
2. Stupp R, Mason WP, Van Den Bent MJ, Weller M, Fisher B, Taphoorn MJ, Belanger K, Brandes AA, Marosi C, Bogdahn U (2005) Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *New England Journal of Medicine* 352: 987-996
3. Gilbert MR, Dignam JJ, Armstrong TS, Wefel JS, Blumenthal DT, Vogelbaum MA, Colman H, Chakravarti A, Pugh S, Won M (2014) A randomized trial of bevacizumab for newly diagnosed glioblastoma. *New England Journal of Medicine* 370: 699-708
4. Gilbert MR, Wang M, Aldape KD, Stupp R, Hegi ME, Jaeckle KA, Armstrong TS, Wefel JS, Won M, Blumenthal DT, Mahajan A, Schultz CJ, Erridge S, Baumert B, Hopkins KI, Tzuk-Shina T, Brown PD, Chakravarti A, Curran WJ, Jr., Mehta MP (2013) Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 31: 4085-4091 doi:10.1200/JCO.2013.49.6968
5. Stupp R, Hegi ME, Gorlia T, Erridge SC, Perry J, Hong Y-K, Aldape KD, Lhermitte B, Pietsch T, Grujicic D (2014) Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071-22072 study): a multicentre, randomised, open-label, phase 3 trial. *The Lancet Oncology* 15: 1100-1108
6. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, Carpentier AF, Hoang-Xuan K, Kavan P, Cernea D (2014) Bevacizumab plus radiotherapy–temozolomide for newly diagnosed glioblastoma. *New England Journal of Medicine* 370: 709-722
7. Wick W, Steinbach JP, Kuker WM, Dichgans J, Bamberg M, Weller M (2004) One week on/one week off: a novel active regimen of temozolomide for recurrent glioblastoma. *Neurology* 62: 2113-2115
8. Brandes AA, Tosoni A, Cavallo G, Bertorelle R, Gioia V, Franceschi E, Biscuola M, Blatt V, Crino L, Ermani M, Gicno (2006) Temozolomide 3 weeks on and 1 week off as first-line therapy for recurrent glioblastoma: phase II study from gruppo italiano cooperativo di neuro-oncologia (GICNO). *Br J Cancer* 95: 1155-1160 doi:10.1038/sj.bjc.6603376
9. Perry JR, Belanger K, Mason WP, Fulton D, Kavan P, Easaw J, Shields C, Kirby S, Macdonald DR, Eisenstat DD, Thiessen B, Forsyth P, Pouliot JF (2010) Phase II trial of continuous dose-intense temozolomide in recurrent malignant glioma: RESCUE study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 28: 2051-2057 doi:10.1200/JCO.2009.26.5520
10. Norden AD, Lesser GJ, Drappatz J, Ligon KL, Hammond SN, Lee EQ, Reardon DR, Fadul CE, Plotkin SR, Batchelor TT, Zhu JJ, Beroukhir R, Muzikansky A, Doherty L, Lafrankie D, Smith K, Tafoya V, Lis R, Stack EC, Rosenfeld MR, Wen PY (2013) Phase 2 study of dose-intense temozolomide in recurrent glioblastoma. *Neuro-oncology* 15: 930-935 doi:10.1093/neuonc/not040
11. Weller M, Tabatabai G, Kastner B, Felsberg J, Steinbach JP, Wick A, Schnell O, Hau P, Herrlinger U, Sabel MC, Wirsching HG, Ketter R, Bahr O, Platten M, Tonn JC, Schlegel U, Marosi C, Goldbrunner R, Stupp R, Homicsko K, Pichler J, Nikkhah G, Meixensberger J, Vajkoczy P, Kollias S, Husing J, Reifenberger G, Wick W, Group DS (2015) MGMT Promoter Methylation Is a Strong Prognostic Biomarker for Benefit from Dose-Intensified Temozolomide Rechallenge in Progressive Glioblastoma: The DIRECTOR Trial. *Clinical cancer research : an official journal of the American Association for Cancer Research* 21: 2057-2064 doi:10.1158/1078-0432.CCR-14-2737
12. Wick W, Puduvalli VK, Chamberlain MC, van den Bent MJ, Carpentier AF, Cher LM, Mason W, Weller M, Hong S, Musib L (2010) Phase III study of enzastaurin compared with

lomustine in the treatment of recurrent intracranial glioblastoma. *Journal of clinical oncology* 28: 1168-1174

13. Batchelor TT, Mulholland P, Neyns B, Nabors LB, Campone M, Wick A, Mason W, Mikkelsen T, Phuphanich S, Ashby LS (2013) Phase III randomized trial comparing the efficacy of cediranib as monotherapy, and in combination with lomustine, versus lomustine alone in patients with recurrent glioblastoma. *Journal of Clinical Oncology* 31: 3212-3218

14. Weller M, van den Bent M, Hopkins K, Tonn JC, Stupp R, Falini A, Cohen-Jonathan-Moyal E, Frappaz D, Henriksson R, Balana C, Chinot O, Ram Z, Reifenberger G, Soffietti R, Wick W, European Association for Neuro-Oncology Task Force on Malignant G (2014) EANO guideline for the diagnosis and treatment of anaplastic gliomas and glioblastoma. *The Lancet Oncology* 15: e395-403 doi:10.1016/S1470-2045(14)70011-7

15. Taal W, Oosterkamp HM, Walenkamp AM, Dubbink HJ, Beerepoot LV, Hanse MC, Buter J, Honkoop AH, Boerman D, de Vos FY, Dinjens WN, Enting RH, Taphoorn MJ, van den Berkmoortel FW, Jansen RL, Brandsma D, Bromberg JE, van Heuvel I, Vernhout RM, van der Holt B, van den Bent MJ (2014) Single-agent bevacizumab or lomustine versus a combination of bevacizumab plus lomustine in patients with recurrent glioblastoma (BELOB trial): a randomised controlled phase 2 trial. *The Lancet Oncology* 15: 943-953 doi:10.1016/S1470-2045(14)70314-6

16. Kreisl TN, Zhang W, Odia Y, Shih JH, Butman JA, Hammoud D, Iwamoto FM, Sul J, Fine HA (2011) A phase II trial of single-agent bevacizumab in patients with recurrent anaplastic glioma. *Neuro-oncology* 13: 1143-1150

17. Vredenburgh JJ, Desjardins A, Herndon JE, 2nd, Dowell JM, Reardon DA, Quinn JA, Rich JN, Sathornsumetee S, Gururangan S, Wagner M, Bigner DD, Friedman AH, Friedman HS (2007) Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clinical cancer research : an official journal of the American Association for Cancer Research* 13: 1253-1259 doi:10.1158/1078-0432.CCR-06-2309

18. Woernle CM, Peus D, Hofer S, Rushing EJ, Held U, Bozinov O, Krayenbuhl N, Weller M, Regli L (2015) Efficacy of Surgery and Further Treatment of Progressive Glioblastoma. *World Neurosurg* doi:10.1016/j.wneu.2015.03.018

19. Franceschi E, Bartolotti M, Tosoni A, Bartolini S, Sturiale C, Fioravanti A, Pozzati E, Galzio R, Talacchi A, Volpin L, Morandi L, Danieli D, Ermani M, Brandes AA (2015) The effect of re-operation on survival in patients with recurrent glioblastoma. *Anticancer research* 35: 1743-1748

20. Vuorinen V, Hinkka S, Farkkila M, Jaaskelainen J (2003) Debulking or biopsy of malignant glioma in elderly people - a randomised study. *Acta neurochirurgica* 145: 5-10 doi:10.1007/s00701-002-1030-6

21. Fogh SE, Andrews DW, Glass J, Curran W, Glass C, Champ C, Evans JJ, Hyslop T, Pequignot E, Downes B, Comber E, Maltenfort M, Dicker AP, Werner-Wasik M (2010) Hypofractionated stereotactic radiation therapy: an effective therapy for recurrent high-grade gliomas. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 28: 3048-3053 doi:10.1200/JCO.2009.25.6941

22. Mizumoto M, Okumura T, Ishikawa E, Yamamoto T, Takano S, Matsumura A, Oshiro Y, Ishikawa H, Sakurai H, Tsuboi K (2013) Reirradiation for recurrent malignant brain tumor with radiotherapy or proton beam therapy. Technical considerations based on experience at a single institution. *Strahlenther Onkol* 189: 656-663 doi:10.1007/s00066-013-0390-6

23. Gutin PH, Iwamoto FM, Beal K, Mohile NA, Karimi S, Hou BL, Lymberis S, Yamada Y, Chang J, Abrey LE (2009) Safety and efficacy of bevacizumab with hypofractionated stereotactic irradiation for recurrent malignant gliomas. *International journal of radiation oncology, biology, physics* 75: 156-163 doi:10.1016/j.ijrobp.2008.10.043

24. Wick W, Fricke H, Junge K, Kobayakov G, Martens T, Heese O, Wiestler B, Schliesser MG, von Deimling A, Pichler J, Vetlova E, Harting I, Debus J, Hartmann C, Kunz C, Platten M, Bendszus M, Combs SE (2014) A phase II, randomized, study of weekly

APG101+reirradiation versus reirradiation in progressive glioblastoma. Clinical cancer research : an official journal of the American Association for Cancer Research 20: 6304-6313 doi:10.1158/1078-0432.CCR-14-0951-T

25. Hundsberger T, Brügge D, Putora PM, Weder P, Weber J, Plasswilm L (2013) Re-irradiation with and without bevacizumab as salvage therapy for recurrent or progressive high-grade gliomas. Journal of neuro-oncology 112: 133-139

26. Weller M, Cloughesy T, Perry JR, Wick W (2013) Standards of care for treatment of recurrent glioblastoma—are we there yet? Neuro-oncology 15: 4-27

27. Putora PM, Oldenburg J (2013) Swarm-based medicine. J Med Internet Res 15: e207 doi:10.2196/jmir.2452

28. Putora P.M. BM, Papachristofilou A., Mariotti F., Paoli B., Plasswilm L. (2010) Dodes (diagnostic nodes) for Guideline Manipulation. J Radiat Oncol Inform 2: 1-8

29. Putora PM, Panje CM, Papachristofilou A, Dal Pra A, Hundsberger T, Plasswilm L (2014) Objective consensus from decision trees. Radiation oncology 9: 270

30. Rothermundt C, Bailey A, Cerbone L, Eisen T, Escudier B, Gillessen S, Grunwald V, Larkin J, McDermott D, Oldenburg J, Porta C, Rini B, Schmidinger M, Sternberg C, Putora PM (2015) Algorithms in the First-Line Treatment of Metastatic Clear Cell Renal Cell Carcinoma-Analysis Using Diagnostic Nodes. The oncologist 20: 1028-1035 doi:10.1634/theoncologist.2015-0145

31. Panje CM, Dal Pra A, Zilli T, D RZ, Papachristofilou A, Herrera FG, Matzinger O, Plasswilm L, Putora PM (2015) Consensus and differences in primary radiotherapy for localized and locally advanced prostate cancer in Switzerland : A survey on patterns of practice. Strahlenther Onkol May 19 doi:10.1007/s00066-015-0849-8

32. Batchelor TT, Mulholland P, Neyns B, Nabors LB, Campone M, Wick A, Mason W, Mikkelsen T, Phuphanich S, Ashby LS, Degroot J, Gattamaneni R, Cher L, Rosenthal M, Payer F, Jurgensmeier JM, Jain RK, Sorensen AG, Xu J, Liu Q, van den Bent M (2013) Phase III randomized trial comparing the efficacy of cediranib as monotherapy, and in combination with lomustine, versus lomustine alone in patients with recurrent glioblastoma. Journal of clinical oncology : official journal of the American Society of Clinical Oncology 31: 3212-3218 doi:10.1200/JCO.2012.47.2464

33. Wick W, Puduvalli VK, Chamberlain MC, van den Bent MJ, Carpentier AF, Cher LM, Mason W, Weller M, Hong S, Musib L, Liepa AM, Thornton DE, Fine HA (2010) Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. Journal of clinical oncology : official journal of the American Society of Clinical Oncology 28: 1168-1174 doi:10.1200/JCO.2009.23.2595

34. Weller M, Cloughesy T, Perry JR, Wick W (2013) Standards of care for treatment of recurrent glioblastoma--are we there yet? Neuro-oncology 15: 4-27 doi:10.1093/neuonc/nos273

35. Mariniello G, Peca C, De Caro Mdel B, Giamundo A, Donzelli R, Maiuri F (2014) Glioblastoma in the elderly: the impact of advanced age on treatment and survival. J Neurol Surg A Cent Eur Neurosurg 75: 276-281 doi:10.1055/s-0033-1349713

36. Kirkpatrick JP, Sampson JH (2014) Recurrent malignant gliomas. Semin Radiat Oncol 24: 289-298 doi:10.1016/j.semradonc.2014.06.006

37. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, Belanger K, Brandes AA, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier A, Lacombe D, Cairncross JG, Eisenhauer E, Mirimanoff RO, European Organisation for R, Treatment of Cancer Brain T, Radiotherapy G, National Cancer Institute of Canada Clinical Trials G (2005) Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. The New England journal of medicine 352: 987-996 doi:10.1056/NEJMoa043330

38. Chinot OL, de La Motte Rouge T, Moore N, Zeaiter A, Das A, Phillips H, Modrusan Z, Cloughesy T (2011) AVAglio: Phase 3 trial of bevacizumab plus temozolomide and

radiotherapy in newly diagnosed glioblastoma multiforme. *Adv Ther* 28: 334-340
doi:10.1007/s12325-011-0007-3

39. Gorlia T, van den Bent MJ, Hegi ME, Mirimanoff RO, Weller M, Cairncross JG, Eisenhauer E, Belanger K, Brandes AA, Allgeier A, Lacombe D, Stupp R (2008) Nomograms for predicting survival of patients with newly diagnosed glioblastoma: prognostic factor analysis of EORTC and NCIC trial 26981-22981/CE.3. *The Lancet Oncology* 9: 29-38
doi:10.1016/S1470-2045(07)70384-4

40. Fiorentino A, Caivano R, Chiumento C, Cozzolino M, Clemente S, Pedicini P, Fusco V (2012) Comorbidity assessment and adjuvant radiochemotherapy in elderly affected by glioblastoma. *Med Oncol* 29: 3467-3471 doi:10.1007/s12032-012-0246-4

41. Balducci M, Fiorentino A, De Bonis P, Chiesa S, Manfreda S, D'Agostino GR, Mantini G, Frascino V, Mattiucci GC, De Bari B, Mangiola A, Micciche F, Gambacorta MA, Colicchio G, Morganti AG, Anile C, Valentini V (2012) Impact of age and co-morbidities in patients with newly diagnosed glioblastoma: a pooled data analysis of three prospective mono-institutional phase II studies. *Med Oncol* 29: 3478-3483 doi:10.1007/s12032-012-0263-3

42. Ackerl M, Flechl B, Dieckmann K, Preusser M, Widhalm G, Sax C, Marosi C (2014) Outcome evaluation in glioblastoma patients older than 65 years: Importance of individual assessment of treatment tolerance. *Clin Neuropathol* 33: 399-406 doi:10.5414/NP300761

43. Bauchet L, Zouaoui S, Darlix A, Menjot de Champfleury N, Ferreira E, Fabbro M, Kerr C, Taillandier L (2014) Assessment and treatment relevance in elderly glioblastoma patients. *Neuro-oncology* 16: 1459-1468 doi:10.1093/neuonc/nou063

44. Gilbert MR, Dignam JJ, Armstrong TS, Wefel JS, Blumenthal DT, Vogelbaum MA, Colman H, Chakravarti A, Pugh S, Won M, Jeraj R, Brown PD, Jaeckle KA, Schiff D, Stieber VW, Brachman DG, Werner-Wasik M, Tremont-Lukats IW, Sulman EP, Aldape KD, Curran WJ, Jr., Mehta MP (2014) A randomized trial of bevacizumab for newly diagnosed glioblastoma. *The New England journal of medicine* 370: 699-708
doi:10.1056/NEJMoa1308573

45. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, Carpentier AF, Hoang-Xuan K, Kavan P, Cernea D, Brandes AA, Hilton M, Abrey L, Cloughesy T (2014) Bevacizumab plus radiotherapy-temozolomide for newly diagnosed glioblastoma. *The New England journal of medicine* 370: 709-722 doi:10.1056/NEJMoa1308345

46. Park JK, Hodges T, Arko L, Shen M, Dello Iacono D, McNabb A, Olsen Bailey N, Kreisl TN, Iwamoto FM, Sul J, Auh S, Park GE, Fine HA, Black PM (2010) Scale to predict survival after surgery for recurrent glioblastoma multiforme. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 28: 3838-3843
doi:10.1200/JCO.2010.30.0582

Fig. 1




Fig. 3

treatments								
centre:	A	B	C	D	E	F	G	H
OP			+	+			+	
SRS			+				+	
BEV	+	+	+	+			+	+
TMZ	+	+	+		+	+	+	+
RT	+	+		+				
LOM				+		+		+
BEV&LOM					+			
OP+TMZ	+	+			+	+		+
OP+BEV&LOM					+			
BSC	+	+			+	+	+	+
OP+LOM	+					+		+
OP+BEV	+	+				+		+
OP+RT&TMZ	+							
OP+RT	+	+						
RT&TMZ	+							
RT&BEV						+		

Fig. 3




Table 1

re-treatment criteria	assessment	simplified criteria
performance status	KPS \leq 50; 60; 70; 90	fit vs. unfit
	ECOG \leq 1	
	good/bad	
age	\leq 50 years; \leq 75 years	
tumour volume	\geq 65 ml; \leq 1.5 cm	operable vs. not operable
rGBM characteristics	new/distant lesion	
	diffuse / local lesion	
	resectable (yes/no)	
	single/ multiple lesions	
	growth kinetics (slow/rapid)	
time from last TMZ treatment	\leq 2;4;6 months	early vs. late
time from last radiotherapy	\leq 6 months	
re-resection	total; subtotal	CRET vs. non-CRET
symptomatic rGBM	yes/no	symptomatic vs. asymptomatic
steroid use	yes/no	
methylated MGMT promoter	yes/no	unchanged

Among 8 neuro-oncology centres 23 re-treatment criteria were disclosed. Based on unification and simplification similar criteria were partially re-named and merged. Abbreviations: KPS, Karnofsky performance status; ECOG, Eastern Cooperative Oncology Group performance status; rGBM, recurrent glioblastoma; CRET, complete resection of contrast enhancing tumour; TMZ, temozolomide; vs., versus